

OCCURRENCE AND CORRELATES OF OVERWEIGHT AND OBESITY AMONG ISLAND PUERTO RICAN YOUTH

Objective and Main Outcome Measures:

This article provides 2005–2008 population-based prevalence data on obesity and overweight among youth residing in Puerto Rico.

Design and Setting: Data for this report are from the Asthma, Depression, and Anxiety in Puerto Rican Youth (ADA) study. Measures included height and weight level data on youth in Puerto Rico aged 10 to 19 years with and without asthma as well as body mass index data on their caregivers.

Participants: A total of 436 youth-caregiver dyads were selected and weighted to represent the general population of youth in Puerto Rico using 2008 US Census data.

Results: Household surveys demonstrated that 40% of youth aged 10 to 19 were overweight or obese. Twenty-five percent met moderate-to-vigorous-intensity physical activity guidelines, however, physical activity was not associated with overweight or obesity in this sample. In multivariate analyses, females were 50% less likely than males to be overweight or obese. Older youth were 73% less likely to be overweight or obese than younger youth. Youth whose parents were obese were more than two times more likely to be overweight or obese than those whose parents were at a desirable weight.

Conclusions: Youth in Puerto Rico have higher rates of overweight and obesity and lower compliance to moderate-to-vigorous-intensity physical activity guidelines than rates reported for youth on the mainland. More population-based research is needed to understand the epidemiology of obesity and overweight among island Puerto Rican youth and the contribution of physical activity to the phenomenon. (*Ethn Dis.* 2011;21(2):163–169)

Key Words: Child, Overweight/epidemiology, Obesity/epidemiology, Physical Activity, Prevalence, Puerto Rico/epidemiology

Jeremiah R. Garza, MPH; Edna Acosta Pérez, PhD, MSc;
Michael Prelep, DPA; William J. McCarthy, PhD;
Jonathan M. Feldman, PhD; Glorisa Canino, PhD;
Alexander N. Ortega, PhD

INTRODUCTION

Obesity has been described as an epidemic because of the relatively high number of individuals who have become overweight or obese over the past 20 years.¹ The rise in obesity is a worldwide phenomenon affecting both children and adults.^{1–2} It is a societal health challenge because excess body weight is the sixth most important risk factor contributing to the overall burden of disease globally, increasing the risk of various chronic diseases.^{2–3}

Childhood and adolescence (collectively, youth) have been proposed as critical periods for the development of this condition.^{2,4} Obesity in early life is of particular concern as approximately one-half of overweight adolescents and over one-third of overweight children remain obese as adults.^{2,5} Childhood obesity is associated with a wide range of serious complications in the short-term, and, if it persists into adulthood, increases the risk of excess illness and premature death.⁶

Angeles (MP) and University of Puerto Rico, Behavioral Sciences Research Institute, Medical Science Campus, San Juan (EAP, GC) and Yeshiva University, Ferkauf Graduate School of Psychology and Albert Einstein College of Medicine, Department of Epidemiology and Population Health, Bronx, New York (JMF).

Address correspondence to Jeremiah R. Garza, MA, MPH; UCLA School of Public Health; 650 Charles E. Young Drive South-Box 951772; Los Angeles, CA 90095; 310.428.1669; 310.206.4453 (fax); J.Garza1@ucla.edu

Generally, higher rates of overweight and obesity are associated with lower physical activity participation.^{7–9} Moreover, a parallel increase in physical inactivity among youth is worrisome, as daily physical activity fosters optimal physical and cognitive development.¹⁰ For these reasons, several countries and their territories view the prevention of obesity as a public health priority that requires a concomitant focus on physical activity.^{2,3}

The US Commonwealth of Puerto Rico, where public health practitioners have been struggling with an increasing prevalence of pediatric obesity and inactivity-related diseases – such as diabetes, high blood pressure and heart disease – is no exception.^{11–14} According to the Centers for Disease Control and Prevention's (CDC) 2005 High School Youth Risk Behavior Survey (YRBS), the most recent data available on overweight/obesity among island Puerto Rican youth, 14% of youth in Puerto Rico are overweight and 12% obese.¹⁵ These island estimates are similar to the 15.8% overweight and 12.0% obese rates published most recently on mainland youth in the 2009 YRBS.¹⁶ Other studies in Puerto Rico that included children recruited from schools and clinics have reported obesity rates as high as 24.6% and 36%, respectively.^{12,13} The 2005 YRBS survey also shows that among those aged ≥ 14 years, fewer Puerto Rican high school students (20.8%) met physical activity guidelines than youth (35.8%) on the US mainland.¹⁵ Accumulation of at least 60 min of

From Department of Health Services, UCLA School of Public Health, Los Angeles (JRG, WJM, ANO) and UCLA Department of Psychology, Los Angeles (WJM) and Department of Community Health Sciences, UCLA School of Public Health, Los

The current literature on obesity among youth in Puerto Rico, including the YRBS, is limited by the use of clinical- or school-based rather than population-based samples, and it has mostly failed to include determinants of obesity, such as level of physical activity and familial or social characteristics.

moderate-to-vigorous-intensity physical activity daily (MVPA) is recommended for youth.^{17–19} Examples of MVPA include moderate-intensity aerobic activity, such as brisk walking, and vigorous-intensity activity, such as running.¹⁹ The lower rates of MVPA in Puerto Rico are troubling given that the US Department of Agriculture (USDA) and the US Department of Health and Human Services recommend regular physical activity to facilitate weight control.^{18,19} While popular media and a small handful of empirical studies have reported on the relationship between obesity and inactivity among youth in Puerto Rico, more population-based research is needed that examines island youth.^{11–14,20–22} The current literature on obesity among youth in Puerto Rico, including the YRBS, is limited by the use of clinical- or school-based rather than population-based samples, and it has mostly failed to include determinants of obesity, such as level of physical activity and familial or social characteristics.

In this study, we sought to determine the strength of the associations between correlates of youth overweight/obesity and physical activity, sociodemographics and parental/caregiver vari-

ables using youth- or parent-reported data from the Asthma, Depression and Anxiety in Puerto Rican Youth (ADA) study.^{23,24} The ADA study also collected simple measures of obesity, as well as physical activity, which allowed us to describe the occurrence and correlates of these variables among youth and their caregivers in the data set. With respect to social variables, while a number of mainland studies have documented higher rates of overweight and obesity among low-socioeconomic groups, there has been a paucity of such studies in Puerto Rico.^{25,26} This study therefore explores the relationship between social variables such as perception of poverty, household income, and parental education, among other variables, with youth overweight and obesity status. This study is unique because it uses a larger, population-based sample compared with prior studies, and it examines the prevalence and correlates of early life (10 to 19 years) overweight/obesity and physical activity on the island.

METHODS

Data for this study are from the third wave (2005–2008) of the ADA study which was specifically designed to assess the associations of asthma and asthma care with child and parental psychiatric disorders among Puerto Rican children aged 4 to 17 years.^{23,24} We were, however, able to collect some limited information on obesity and physical activity in this third wave. Wave one and two did not include these questions; thus, the analyses presented here involve cross-sectional data from wave three. Wave one was conducted from 1999–2000, and wave two was a one-year follow-up from 2000–2001. Details regarding the sampling design and procedures have been previously described for wave one and two.^{23,24} The sample and measures are described briefly here, with particular focus paid

to describing wave three of the study, which has not been previously reported.

Youth aged 4 to 17 years living on the island of Puerto Rico made up an island-wide household probability sample stratified by four dimensions: urban vs rural areas, Puerto Rico's health reform areas, child's age, and sex. A total of 2,102 children from the community were deemed eligible. At wave one, 1,886 children and their caregivers were interviewed for a response rate of 90.1%. A total of 1,789 caregiver-youth dyads from wave one were interviewed at wave two, for a 94.9% retention rate at the one-year follow-up.

For wave three, we used direct mail to recruit participants from wave two. The goal for the ADA study was to obtain a representative community sample including youth and young adults stratified into four groups (asthma and anxiety/depression; asthma no anxiety/depression; anxiety/depression no asthma; neither asthma nor anxiety/depression). Using simple random selection, 825 households were contacted, from which 656 youth and young adults, aged 10 to 25 years, were interviewed for a response rate of 79.5%. Because the current study focuses only on youth and not young adults (because some of the youth in wave 1 were young adults by wave three), we included caregiver-youth dyads with male and female youth aged 10 to 19 years ($n=436$) at the time of wave three data collection.^{23,24}

Blinded interviewers conducted interviews in the families' homes and different interviewers were used for the youth/young adult and caregiver interviews. The adult informant was the participant's biological mother for ~89% of the interviews. All interviews were audiotaped, and 15% were randomly reviewed for quality control. The study protocol was approved by the institutional review boards of the University of Puerto Rico, Medical Sciences Campus and the University of California Los Angeles. Caregiver

consent and child assent were obtained for youth aged <18. Consent was obtained for participants ≥ 18 years. In order for a youth/young adult to participate, caregivers were also required to participate in the study to provide information about themselves and their progeny.

The survey collected demographic information, body mass index (BMI), and physical activity level, among other measures.^{23,24,27-29} Parent-reported demographic variables included parental education, marital status, work status, household income, household composition, perception of poverty, and child's age and sex.³¹ Body mass index was based on child height and weight information obtained from parental report for youth <17 years. Youth aged 17 to 19 years provided information on their own weight and height. For youth aged <20, the 85th percentile for age- and sex-specific BMI levels using CDC growth chart norms was used as the cutpoint for child classification as overweight and the 95th percentile for classification as obese.²⁷ All youth weight below the 85th percentile was termed, desirable weight. Weight status was interpreted for caregivers (≥ 20 years) using CDC-defined standard weight status categories (ie, desirable weight, overweight, obese).^{27,28} We used a measure of youth compliance to the federal recommendation of at least 60 minutes of MVPA daily.¹⁸ A two-item PACE+ Adolescent Physical Activity Measure assessed the number of days youth had accumulated at least 60 minutes of MVPA per day during the past seven days and for a typical week.²⁹ Information regarding MVPA was obtained by parental report for children <17 years, while youth 17 to 19 years provided their own information. We report a composite average of the two items, yielding a score of the number of days per week during which the youth accumulated 60 minutes of MVPA.²⁹ Five or more days per week met the federal guideline for youth.²⁹

Analyses were weighted to account for the complex sampling design, to correct for differential nonresponse, and to represent the general population of youth in Puerto Rico using 2008 US Census data. The estimation of design weights used to make our sample representative of youth in Puerto Rico was accomplished in two stages. We estimated the participants' probability of selection during the third wave and made an additional adjustment for the response rate. The probability of selection took into account that for wave three we selected a different number of participants from four strata of different sizes. The inverse of this final probability was used to estimate the initial design weights. The design weight estimated during this first stage made our sample representative of the youth population in Puerto Rico in the year 2000 using 2000 US Census Data. In the second stage we made an additional adjustment to our design weights by post-stratifying the data to the population of youth in Puerto Rico as documented in 2008 US Census data. The results were estimated with SUDAAN 10 software to adjust standard errors for multistage sampling, with youth-caregiver dyads nested within households and households nested within primary sampling units.³⁰

Chi-square tests and logistic regression models were used to examine associations among youth overweight/obesity with physical activity, socioeconomic status, parent marital status and parent BMI.

RESULTS

Of the entire youth sample, 17.9% were reported as overweight and 21.5% were reported as obese. According to the unadjusted results in Table 1, the odds of youth aged 15 to 19 years being overweight/obese were 35% (OR=.35, 95% CI=.22, .58) of the odds of youth aged 10 to 14 years. Female youth were

37% (OR=.63, 95% CI=.38, 1.02) less likely to be overweight/obese than males. Youth whose parents reported an annual household income of $\leq \$6,000$ were 2.06 (95% CI=.96, 4.42) times more likely to be overweight/obese than those from households with annual family incomes over \$25,000.

In terms of parents' demographic characteristics, 67% of the sample were married; over 70% were high school or college educated and 55% perceived their family as living well financially. Overweight and obese caregivers made up 35.0% and 36.3%, respectively, of the sample. Obese caregivers, compared to desirable weight parents, were more likely (OR=2.69, 95% CI=1.34, 5.37) to have overweight or obese children. Approximately 25% of youth complied with the federal MVPA recommendation, while 36% of parents reported their children as meeting the recommendation.

Table 2 shows the adjusted results for the associations involving demographic factors and physical activity with youth overweight/obese status. Older youth had lower odds (OR=0.27, 95% CI=.16, .47) of being overweight or obese than younger youth. Female youth had lower odds (OR=.50, 95% CI=.28, .91) of being overweight or obese than male youth. Youth reporting more days per week of at least 60 minutes of exercise were moderately less likely to be overweight or obese (OR=.96, 95% CI=.88, 1.05). Obese caregivers, compared to desirable weight parents, were more likely (OR=2.78, 95% CI=1.38, 5.61) to have overweight or obese children.

CONCLUSION

Our prevalence estimates of overweight and obesity among island Puerto Rican youth differ from previous published studies. For example, the CDC's 2005 YRBS, which used a school-based

Table 1. Unadjusted associations of youth and family/parental characteristics with youth overweight/obesity status

| Characteristics | N=436 | | Overweight/obese (n=152) | Desirable weight (not overweight/not obese) (n=227) | OR (95% CI) |
|---|-------|-----------|-----------------------------|---|------------------|
| | n | % or mean | % or mean (SE) | % or mean (SE) | |
| Youth Characteristics | | | | | |
| Age Category | | | | | |
| 10 to 14 years | 157 | 47.25 | 52.26 (4.97) | 47.74 (4.97) | reference |
| 15 to 19 years | 222 | 52.75 | 27.98 (3.38) | 72.02 (3.38) | .35 (.22–.58) |
| Male | 196 | 50.59 | 44.96 (4.42) | 55.04 (4.42) | reference |
| Female | 183 | 49.41 | 33.81 (4.10) | 66.19 (4.10) | .63 (.38–1.02) |
| Physical activity (number of days in usual week exercised 60+ minutes) | 376 | 3.46 | 3.46 (.23) | 3.46 (.21) | 1.00 (.92–1.09) |
| Smoke Exposure | | | | | |
| No | 238 | 68.47 | 39.16 (3.66) | 60.84 (3.66) | reference |
| Yes | 132 | 31.53 | 40.88 (5.57) | 59.12 (5.57) | 1.07 (.64–1.82) |
| Youth Self-Report PACE+ Adolescent Physical Activity Measure | | | | | |
| Not met moderate to vigorous physical activity guidelines | 283 | 74.72 | 40.30 (3.60) | 59.70 (3.60) | reference |
| Met moderate to vigorous physical activity guidelines | 92 | 25.28 | 35.73 (5.39) | 64.27 (5.39) | .82 (.50–1.37) |
| Caregiver* Report PACE+ Adolescent Physical Activity Measure | | | | | |
| Not met moderate to vigorous physical activity guidelines | 235 | 63.94 | 39.12 (3.81) | 60.88 (3.81) | reference |
| Met moderate to vigorous physical activity guidelines | 138 | 36.06 | 40.44 (5.00) | 59.56 (5.00) | 1.06 (.65–1.71) |
| Family/Parental Characteristics | | | | | |
| Household composition (number of people in household) | 379 | 4.17 | 4.21 (.12) | 4.14 (.08) | 1.05 (.88–1.25) |
| Maternal figure education | | | | | |
| Less than high school | 60 | 15.27 | 35.13 (7.44) | 64.87 (7.44) | .83 (.40–1.69) |
| High school | 120 | 31.12 | 40.79 (5.71) | 59.21 (5.71) | 1.05 (.59–1.86) |
| Some college | 194 | 53.62 | 39.63 (4.21) | 60.37 (4.21) | reference |
| Paternal figure education | | | | | |
| Less than high school | 74 | 28.24 | 38.00 (7.18) | 62.00 (7.18) | .96 (.46–1.99) |
| High school | 87 | 31.28 | 38.42 (6.50) | 61.58 (6.50) | .97 (.47–2.04) |
| Some college | 96 | 40.49 | 39.05 (5.80) | 60.95 (5.80) | reference |
| Income | | | | | |
| ≤\$6,000 | 77 | 20.21 | 51.99 (6.97) | 48.01 (6.97) | 2.06 (.96–4.42) |
| \$ 6,001–\$12,000 | 81 | 23.63 | 33.93 (5.76) | 66.07 (5.76) | .98 (.49–1.94) |
| \$12,001–\$25,000 | 98 | 27.78 | 38.71 (5.44) | 61.29 (5.44) | 1.20 (.62–2.34) |
| >\$25,000 | 94 | 28.39 | 34.45 (5.75) | 65.55 (5.75) | reference |
| Perception of poverty | | | | | |
| Live poorly | 40 | 7.66 | 33.41 (7.69) | 66.59 (7.69) | .64 (.30–1.39) |
| Live check to check | 140 | 37.81 | 34.00 (4.52) | 66.00 (4.52) | .66 (.39–1.12) |
| Live well | 197 | 54.53 | 43.89 (4.55) | 56.11 (4.55) | reference |
| Employment (maternal figure) | | | | | |
| No | 170 | 43.50 | 42.92 (4.42) | 57.08 (4.42) | reference |
| Yes | 204 | 56.50 | 36.52 (4.05) | 63.48 (4.05) | .77 (.48–1.22) |
| Employment (paternal figure) | | | | | |
| No | 65 | 27.69 | 36.57 (7.22) | 63.43 (7.22) | reference |
| Yes | 192 | 72.31 | 39.32 (4.38) | 60.68 (4.38) | 1.12 (.57–2.24) |
| Caregiver marital status | | | | | |
| Married/living with couple | 255 | 67.13 | 38.45 (3.86) | 61.55 (3.86) | reference |
| Separated/divorced/widowed | 101 | 27.70 | 41.80 (5.99) | 58.20 (5.99) | 1.15 (.64–2.07) |
| Never married | 21 | 5.17 | 37.86 (12.65) | 62.14 (12.65) | .98 (.33–2.91) |
| Caregiver BMI weight status categories* | | | | | |
| Underweight | 5 | 1.37 | 56.75 (26.32) | 43.25 (26.32) | 3.11 (.34–28.36) |
| Normal weight | 101 | 27.67 | 29.69 (6.03) | 70.31 (6.03) | reference |
| Overweight | 128 | 34.67 | 33.20 (4.41) | 66.80 (4.41) | 1.18 (.59–2.34) |
| Obese | 134 | 36.30 | 53.15 (5.21) | 46.85 (5.21) | 2.69 (1.34–5.37) |

* Primary caregiver could be the mother or father, but mostly they were mothers (~89%)

Table 2. Multivariate associations of age, sex, income, physical activity and caregiver BMI with youth overweight/obesity status*

| Variables | OR (95% CI) |
|--|------------------|
| Age Category | |
| 10 to 14 years | reference |
| 15 to 19 years | .27 (.16-.47) |
| Sex | |
| Male | reference |
| Female | .50 (.28-.91) |
| Income | |
| ≤\$6,000 | 1.70 (.76-3.80) |
| \$6,001-\$12,000 | .90 (.44-1.83) |
| \$12,001-\$25,000 | .97 (.44-2.14) |
| >\$25,000 | reference |
| Physical Activity (number of days in usual week exercise 60 minutes) | .96 (.88-1.05) |
| Caregiver BMI weight status categories† | |
| Underweight | 3.48 (.36-33.67) |
| Normal weight | reference |
| Overweight | 1.43 (.70-2.91) |
| Obese | 2.78 (1.38-5.61) |

* For this regression, we used 341 cases out of a total of 436; missing data caused 105 to be dropped through listwise deletion
 † Primary caregiver could be the mother or father, but mostly they were mothers (~89%)

sample, provides the most current estimates available for the island. It found that 14% and 12% of Puerto Rican 9th through 12th graders were overweight and obese, respectively, compared with 15.7% and 13.1% for US mainland youth.¹⁵ Compared with the latest estimates of the general US mainland youth, our estimates are higher; they are higher than the 15.8% overweight and 12.0% obese reported among high school youth by the CDC's 2009 YRBS and the 18.1% obese reported by the 2007-2008 National Health and Nutrition Examination Survey.^{16,32} With respect to youth physical activity, our estimates of the percentage of Puerto Rican youth meeting federally recommended MVPA guidelines were higher than the 20.8% reported for Puerto Rican island youth in the 2005 YRBS.¹⁵

Regarding our estimates of overweight/obesity, in comparison, a study of US Latino children seen at community health centers in medically underserved areas (including Puerto Rico) reported an obesity estimate of 24.6%.¹³ A separate study of 158

children receiving pediatric care at the San Juan City Hospital and a primary care clinic in Puerto Rico reported an obesity prevalence rate of 36%.¹² Two possible reasons for these differences in findings are: 1) school samples are skewed towards nonrepresentation of older adolescents in places such as Puerto Rico where school truancy is high; and 2) some of the studies included clinical samples with characteristics that might place them at higher risk for being overweight or obese compared to our broader, more representative sample.

The fact that older youth in our study exhibited lower odds of being overweight or obese than younger youth indicates a significant inverse effect of age. This finding is inconsistent with the current literature which shows that overweight children are more likely to become overweight in adolescence and adulthood than before adolescence.^{1-5,33} Our observation may be partially explained by the common growth spurt that occurs during the adolescent years.^{34,35} A good fraction of youth in our sample are experiencing such growth

spurts, and to the extent that their height is increasing faster than their weight, this would help to explain the decline in obesity risk with increasing age in this population as measured by BMI. This may merely demonstrate a temporary artifact of how youth physical development affects BMI and may not be related to lifestyle choices known to affect weight long term.

The lower odds of Puerto Rican female youth in our study for being overweight/obese compared with male youth is consistent with national data on sex differences among Latinos and stand in contrast to the sex differences among African American adolescents, where girls are at higher risk than boys.³⁶ Further study of sex differences in weight status among Puerto Rican youth is important because the causes of overweight/obesity may differ in girls and boys and be mediated by ethnicity.^{36,37}

We did not find a significant relationship between social variables such as perception of poverty, household income, and parental education, with youth overweight and obesity. While it has been well-documented that low-socioeconomic-status groups are disproportionately affected by obesity at all ages,^{25,26} our null findings may be an artifact of examining these relationships in an island population that is generally low-income (45% of the population live below the poverty level);³⁸ over 70% of our sample reported annual household incomes of ≤\$25,000.

The ~70% of parents who were overweight or obese in our study is higher than previously reported prevalence estimates for adults in Puerto Rico.²² The observation that overweight or obese parents were more likely to have overweight or obese children is consistent with studies showing that young children with at least one obese parent have greater odds for becoming obese themselves than children with desirable weight parents.³⁹⁻⁴⁵

In sum, a greater percentage of island youth are overweight and obese compared with corresponding estimates for US mainland youth.

In sum, a greater percentage of island youth are overweight and obese compared with corresponding estimates for US mainland youth. We also report lower Puerto Rican youth compliance to federal physical activity guidelines than is the case for mainland youth. Our findings also suggest a trend toward more male youth being overweight/obese than female youth. Children of obese parents were more likely to be overweight/obese themselves than those with desirable weight parents. Developers of policies and programs designed to prevent or reduce youth obesity in Puerto Rico would do well to take these correlates of obesity into account in design interventions tailored for specific subgroups and in designing future observational research.

LIMITATIONS

There are limitations that should be considered when interpreting these results. First, our findings are based on self-report rather than objective measures of height, weight, and physical activity. Studies have shown that parental reported weight and height may underestimate true weight among girls and overestimate height in boys.⁴⁶ Although typically obesity prevalence estimates derived from self-report are likely to be lower than they would be were objective measures available, they are nonetheless useful for tracking trends over time and for comparing island estimates to comparable mainland estimates.⁴⁷ Further, BMI-for-age

percentile (based on self-reported height and weight) is a proxy measure of weight status, correlates with body fat, and is recommended for assessing weight status in youth aged 2–20.^{48,49} Similarly, while it is not ideal to base physical activity prevalence estimates on self-report measures because of the potential for inaccuracy and bias, self-report was the only feasible validated approach for the present study.⁵⁰ The composite physical activity measure employed has been shown previously to provide a reliable estimate of adolescents' physical activity behavior and to correlate significantly with an objective measure of physical activity.^{29,50} Finally, it is important to note that wave three of the ADA study stratified the selection of subjects based on the youth's asthma and anxiety and depression statuses in wave two, however, we accounted for this stratification in the analyses using sampling weights. Given the limited state of the literature on this topic within the population, we believe that our new weighted, population-based data provide insight into the occurrence of obesity and MVPA among Puerto Rican island youth.

ACKNOWLEDGMENTS

The authors thank Dr Carlos Toro for analyzing our census sample data and Pedro García and Dr. Rafael Ramirez for conducting the statistical analyses. None of the authors have a conflict of interest. This study was supported by grant R01 MH069849 funded by National Institute of Mental Health (NIMH), R25 RR17589 funded by the National Center for Research Resources (NCRR), P50 HL105188 funded by the National Heart, Lung and Blood Institute (NHLBI) and 5P60 MD002261-02 funded by the National Center on Minority Health and Health Disparities (NCMHHHD).

REFERENCES

1. Bray GA, Bellanger T. Epidemiology, trends, and morbidities of obesity and the metabolic syndrome. *Endocrine*. 2006;29(1):109–117.
2. Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes*. 2006;1(1):11–25.

3. Haslam DW, James WPT. Obesity. *Lancet*. 2005;366(9492):1197–1209.
4. Dietz WH. Critical periods in childhood for the development of obesity. *Am J Clin Nutr*. 1994;59(5):955–959.
5. Power C, Lake JK, Cole TJ. Measurement and long-term health risks of child and adolescent fatness. *Int J Obes Relat Metab Disord*. 1997;21(7):507–526.
6. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002;360(9331):473–482.
7. Eaton DK, Kann L, Kinchen S, et al. Youth Risk Behavior Surveillance—United States. *J Sch Health*. 2006;76(7):353–372.
8. Brodersen NH, Steptoe A, Boniface DR, Wardle J. Trends in physical activity and sedentary behavior in adolescence: ethnic and socioeconomic differences. *Br J Sports Med*. 2007;41(3):140–144.
9. Whit-Glover MC, Taylor WC, Floyd MF, Yore MM, Yancey AK, Matthews CE. Disparities in physical activity and sedentary behaviors among US children and adolescents: prevalence, correlates, and intervention implications. *J Public Health Policy*. 2009;30:309–334.
10. Roberts CK, Freed BA, McCarthy WJ. Low aerobic fitness and obesity are associated with lower standardized test scores in children. *J Pediatr*. 2010;156(5):711–718.
11. Associated Press. Big trouble in little Puerto Rico: Obese kids - Governor preparing to declare childhood obesity an island-wide emergency. 2007. Available at <http://www.msnbc.msn.com/id/18768818/>. Last accessed October 12, 2009.
12. Otero-González M, García-Fragoso L. Prevalence of overweight and obesity in a group of children between the ages of 2 to 12 years old in Puerto Rico. *P R Health Sci J*. 2008;27(2):159–161.
13. Stettler N, Elliot MR, Kallan MJ, Auerbach SB, Kumanyika SK. High prevalence of overweight among pediatric users of community health centers. *Pediatrics*. 2005;116(3):381–388.
14. Venegas HL, Pérez CM, Suárez EL, Guzmán M. Prevalence of obesity and its association with blood pressure, serum lipids, and selected lifestyles in a Puerto Rican population of adolescents 12–16 years of age. *P R Health Sci J*. 2003;22(2):137–143.
15. Centers for Disease Control and Prevention. Youth Risk Behavior Survey. Puerto Rico vs. United States Physical Activity. 2005. Available at <http://www.cdc.gov/healthyyouth/yrbs/index.htm>. Last accessed October 12, 2009.
16. Centers for Disease Control and Prevention. Youth Risk Behavior Survey. 2009. Available at: <http://apps.nccd.cdc.gov/youthonline/App/>

- Default.aspx?SID=HS. Last accessed December 23, 2010.
17. Strong WB, Malina RM, Blimkie CJ, et al. Evidence based physical activity for school-age youth. *J Pediatr*. 2005;146(6):732-737.
 18. U.S. Department of Agriculture (USDA). *Dietary Guidelines for Americans* USDA, 2005;42.
 19. U.S. Department of Health and Human Services. Physical Activity Guidelines for Americans. Available at <http://www.health.gov/paguidelines/pdf/paguide.pdf>. Last accessed: November 21, 2009.
 20. Estadísticas de Salud en Puerto Rico (Puerto Rico Health Statistics). Tendenciaspr.com. Available at http://www.tendenciaspr.com/Salud/Salud.html#anchor_319. Last accessed: December 10, 2009.
 21. Ho GYF, Qian H, Kim MY, et al. Health disparities between island and mainland Puerto Ricans. *Rev Panam Salud Publica*. 2006;19(5):331-339.
 22. Centers for Disease Control and Prevention. Behavior Risk Factor Surveillance System. 2009. Available at: <http://www.cdc.gov/BRFSS/>. Last accessed December 23, 2010.
 23. Canino G, Shrout PE, Rubio-Stipec M, et al. The DSM-IV rates of child and adolescent disorders in Puerto Rico: prevalence, correlates, service use, and the effects of impairment. *Arch Gen Psychiatry*. 2004;61(1):85-93.
 24. Feldman JM, Ortega AN, McQuaid EL, Canino G. Comorbidity between asthma attacks and internalizing disorders among Puerto Rican children at one-year follow-up. *Psychosomatics*. 2006;47(4):333-339.
 25. Miech RA, Kumanyika SK, Stettler N, Link BG, Phelan JC, Chang VW. Trends in the association of poverty with overweight among US adolescents, 1971-2004. *JAMA*. 2006;295(20):2385-2393.
 26. Wang Y, Beydoun MA. The obesity epidemic in the United States - gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev*. 2007;29:6-28.
 27. Centers for Disease Control and Prevention. Healthy Weight - it's not a diet, it's a lifestyle! Body Mass Index. Available at <http://www.cdc.gov/healthyweight/assessing/bmi/>. Last accessed: November 1, 2009.
 28. Centers for Disease Control and Prevention. Growth Charts. Available at http://www.cdc.gov/growthcharts/percentile_data_files.htm. Last accessed: January 14, 2010.
 29. Prochaska JJ, Sallis JF, Long B. A physical activity screening measure for use with adolescents in primary care. *Arch Pediatr Adolesc Med*. 2001;155(5):554-559.
 30. Research Triangle Institute: SUDDAN Software for the Statistical Analysis of Correlated Data. Research Triangle Park, N.C., 2002.
 31. Gore S, Aseltine RH Jr, Colton ME. Social structure, life stress and depressive symptoms in a high school-aged population. *J Health Soc Behav*. 1992;33(2):97-113.
 32. Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high body mass index in US children and adolescents, 2007-2008. *JAMA*. 2010;303(3):242-249.
 33. Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T. Do obese children become obese adults? *Prev Med*. 1993;22(2):167-177.
 34. Kuczumski RJ, Ogden CL, Guo SS, et al. 2000 CDC growth charts for the United States: methods and development. National Center for Health Statistics. *Vital Health Stat*. 2002;11(246)
 35. Ogden CL, Kuczumski RJ, Flegal KM, et al. Centers for Disease Control and Prevention 2000 Growth Charts for the United States: improvements to the 1977 National Center for Health Statistics version. *Pediatrics*. 2002;109(1):45-60.
 36. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*. 2006;295(13):1549-1555.
 37. Bjornelv S, Lydersen S, Holmen J, et al. Sex differences in time trends for overweight and obesity in adolescents: the Young-HUNT study. *Scand J Public Health*. 2009;37(8):881-889.
 38. Welcome to Puerto Rico. [welcome.topuertorico.org](http://www.topuertorico.org). Available at <http://www.topuertorico.org/economy.shtml>. Last accessed: February 2, 2011.
 39. Lake JK, Power C, Cole TJ. Child to adult body mass index in the 1958 British birth cohort: associations with parental obesity. *Arch Dis Child*. 1997;77(5):376-381.
 40. Davey Smith G, Steer C, Leary S, Ness A. Is there an intrauterine influence on obesity? evidence from parent child associations in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Arch Dis Child*. 2007;92(10):876-880.
 41. Lawlor DA, Timpson NJ, Harbord RM, et al. Exploring the developmental overnutrition hypothesis using parental-offspring associations and FTO as an instrumental variable. *PLoS Med*. 2008;5(3):e33.
 42. Danielzik S, Langnase K, Mast M, Spethmann C, Muller MJ. Impact of parental BMI on the manifestation of overweight 5-7 year old children. *Eur J Nutr*. 2002;41(3):132-138.
 43. Lawlor DA, Smith GD, O'Callaghan MJ, et al. Epidemiologic evidence for the fetal overnutrition hypothesis: findings from the mater-university study of pregnancy and its outcomes. *Am J Epidemiol*. 2007;165(4):418-424.
 44. Kivimaki M, Lawlor DA, Smith GD, et al. Substantial intergenerational increases in body mass index are not explained by the fetal overnutrition hypothesis: the cardiovascular risk in young Finns study. *Am J Clin Nutr*. 2007;86(5):1509-1514.
 45. Sekine M, Yamagami T, Hamanishi S, et al. Parental obesity, lifestyle factors and obesity in preschool children: results of the Toyama Birth Cohort study. *J Epidemiol*. 2002;12(1):33-39.
 46. Dhaliwal SS, Howat P, Bejoy T, Welborn TA. Self-reported weight and height for evaluating obesity control programs. *Am J Health Behav*. 2010;34(4):489-499.
 47. Sherry B, Jefferds ME, Grummer-Strawn LM. Accuracy of adolescent self-report of height and weight in assessing overweight status: a literature review. *Arch Pediatr Adolesc Med*. 2007;161(12):1154-1161.
 48. Krebs NF, Himes JH, Jacobson D, et al. Assessment of child and adolescent overweight and obesity. *Pediatrics*. 2007;120(4):193-228.
 49. Mei Z, Grummer-Strawn LM, Pietrobelli A, Goulding A, Goran MI, Dietz WH. Validity of body mass index compared with other body-composition screening indexes for assessment of body fatness in children and adolescents. *Am J Clin Nutr*. 2002;75(6):978-985.
 50. Sallis JF, Taylor WC, Dowda M, Freedson PS, Pate RR. Correlates of vigorous physical activity for children in grades 1 through 12: comparing parent-reported and objectively measured physical activity. *Pediatr Exerc Sci*. 2002;14(1):30-44.

AUTHOR CONTRIBUTIONS

Design concept of study: Garza, Prelip, Feldman, Canino, Ortega
Acquisition of data: Garza, Acosta Pérez, Canino, Ortega
Data analysis and interpretation: Garza, Acosta Pérez, Prelip, McCarthy, Canino, Ortega
Manuscript draft: Garza, Acosta Pérez, Prelip, McCarthy, Feldman, Canino, Ortega
Statistical expertise: Garza, McCarthy, Ortega
Acquisition of funding: Garza, Canino, Ortega
Administrative: Garza, Prelip, Feldman, Ortega
Supervision: Garza, Acosta Pérez, Canino, Ortega